

Altogen Biosystems

11200 Manchaca Rd, Ste 203

Austin, TX 78748 USA

Tel. (702) 349-6103

E-mail: techserv@altogen.com

Web: altogen.com

D407 Transfection Protocol (Kit for Retinal Pigment Epithelial Cells)

Catalog No. 1743 Size: 0.5 ml

Catalog No. 1744 Size: 1.5 ml

Catalog No. 7033 Size: 8.0 ml

Contents and Shipping:

D407 Transfection Kit includes D407 Transfection Reagent (0.5 ml / 1.5 ml / 8.0 ml), Transfection Enhancer (0.5 ml), and Complex Condenser (0.5 ml). D407 Transfection Reagent is supplied in liquid form at a concentration of 0.25 mg/ml, shipped at ambient temperature.

Description:

Cationic lipid based reagent is a proprietary formulation optimized for transfection of DNA and RNA into D407 cells.

Product Qualification:

D407 Transfection Reagent is tested functionally by transfection of D407 cells with a small interfering RNAs targeting 3 different genes (Lamin A/C, GAPDH, Cyclophilin B). Transfection Reagent is tested for absence of nuclease contamination and microbial contamination.

Storage:

Store reagent at 4°C upon receipt. If stored properly, reagent is stable for 12 months.

Intended Use:

For *in vitro* use only.

Safety Data Sheet (SDS):

SDS documents are available online at altogen.com

In Vivo Transfection Kits (for compound testing in rodents):

- Catalog #5010 / 5011 / 5012 - Lipid *In Vivo* Transfection Kit
- Catalog #5020 / 5021 / 5022 - Polymer *In Vivo* Transfection Kit
- Catalog #5030 / 5031 / 5032 - Nanoparticle *In Vivo* Transfection Kit
- Catalog #5040 / 5041 / 5042 - PEG-Liposome *In Vivo* Transfection Kit
- Catalog #5050 / 5051 / 5052 - Pancreas *In Vivo* Transfection Kit
- Catalog #5060 / 5061 / 5062 - Liver *In Vivo* Transfection Kit
- Catalog #5070 / 5071 / 5072 - Kidney *In Vivo* Transfection Kit

Transfection Controls and Recommended Products:

- Catalog #4060 - GFP-expressing plasmid DNA (25 ug)
- Catalog #4061 - Cell Cycle Arrest siRNA (5 nmol)
- Catalog #4062 - Apoptosis Inducing siRNA (5 nmol)

To Place an Order:

Both domestic (USA) and international orders can be placed online at altogen.com

Purchase Order (PO) should be e-mailed to orders@altogen.com

Transfection Resource: altogen.com/transfection-resource

Altogen Labs. GLP Compliant Pre-clinical CRO Laboratory Services: AltogenLabs.com

Recommended Transfection Protocols (for 24-well plate):

| D407 Standard Transfection Protocol (24-well plate): | D407 Reverse Transfection Protocol (24-well plate): |
|---|---|
| <ol style="list-style-type: none"> 1. Plate 7,500 - 12,000 D407 cells per well in 0.5 ml of complete growth medium 12–24 hours prior to transfection 2. Wash with 1xPBS and add 0.5 ml of fresh growth medium 3. Prepare transfection complexes by mixing 40 µl of serum-free medium, 5.5 µl of transfection reagent, and <ul style="list-style-type: none"> • 750 ng DNA (or mRNA), or • 30 nM - 50 nM of siRNA (or microRNA) <i>*Referred to a final volume including growth medium</i> 4. Incubate transfection complexes at RT for 15 - 30 minutes 5. <u>Optional</u>: Add 2 µl of Complex Condenser. This reagent reduces the size of transfection complex, therefore increasing transfection efficiency; however it may increase cell toxicity 6. Add prepared transfection complexes to 0.5 ml of complete growth medium with D407 cells (from step 2) 7. Incubate cells at 37°C in a humidified CO₂ incubator 8. Assay for phenotype or target gene expression 48 - 72 hours after transfection | <ol style="list-style-type: none"> 1. Prepare D407 cell suspension: <ol style="list-style-type: none"> a. Trypsinize cells (0.05% Trypsin) for 3-5 minutes at 37°C b. Dilute in complete growth medium to 5 x 10⁴ cells/ml 2. Prepare transfection complexes by mixing 40 µl of serum-free medium, 5.5 µl of transfection reagent, and <ul style="list-style-type: none"> • 750 ng DNA (or mRNA), or • 30 nM - 50 nM of siRNA (or microRNA) <i>*Referred to a final volume including growth medium</i> 3. Incubate transfection complexes at RT for 15 - 30 minutes 4. <u>Optional</u>: Add 2 µl of Complex Condenser. This reagent reduces the size of transfection complex, therefore increasing transfection efficiency; however it may increase cell toxicity 5. Plate 15,000 - 25,000 cells per well in 0.5 ml of complete growth medium (from step #1) into culture plate 6. Add prepared transfection complexes (from step 3 or 4) 7. Incubate cells at 37°C in a humidified CO₂ incubator 8. Assay for phenotype or target gene expression 48 - 72 hours after transfection |
| <u>Optional</u> : Transfection efficiency can be increased by addition of Transfection Enhancer reagent. Add 2 µl of Transfection Enhancer reagent 12-24 hours after transfection | <u>Optional</u> : Transfection efficiency can be increased by addition of Transfection Enhancer reagent. Add 2 µl of Transfection Enhancer reagent 12-24 hours after transfection |
| If the viability of D407 cells being transfected is affected at 16 - 24 hours post-transfection, the level of cytotoxicity can be decreased by changing the growth medium and eliminating redundant exposure of cells to transfectant | If the viability of D407 cells being transfected is affected at 16 - 24 hours post-transfection, the level of cytotoxicity can be decreased by changing the growth medium and eliminating redundant exposure of cells to transfectant |

Scaling Up or Down Transfections:

| Culture Vessel Surface Area (cm ²) | Volume of Growth Medium (ml) | Transfection Reagent (µl) | Complex Condenser (µl) | Transfection Enhancer (µl) |
|--|------------------------------|---------------------------|------------------------|----------------------------|
| 96-well, 0.3 cm x cm | 0.12 | 1.5 | 0.3 | 0.3 |
| 24-well, 2 cm x cm | 0.5 | 5.5 | 2 | 2 |
| 12-well, 4 cm x cm | 1 | 12 | 4 | 4 |
| 6-well, 10 cm x cm | 3 | 35 | 12 | 12 |
| 60-mm, 20 cm x cm | 5 | 60 | 20 | 20 |
| 10-cm, 60 cm x cm | 15 | 180 | 60 | 60 |

Optimizing Transfection:

To obtain the highest transfection efficiency, optimize transfection conditions by varying D407 cell density and amount of transfection reagent. High passage of D407 cells and use of antibiotics (or growth factors) may require using larger volumes of D407 transfection reagent per reaction.

Certificate of Analysis:

Transfection reagent was tested *in vitro* and passed QC. Transfection reagent was tested for absence of nuclease contamination and microbial contamination. Rev. 04/29/2025.

Limited Use Label License:

The purchase of this product conveys to the purchaser the limited right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. This product is for research purposes only and is not for use in commercial applications of any kind. For information on obtaining additional rights, please contact Altogen Biosystems at orders@altogen.com.

Safety Data Sheet:

Please refer to altogen.com for Safety Data Sheet of the product.

Limited Product Warranty:

This warranty limits our liability to replacement of this product. No other warranties of any kind, express or implied, including, without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Altogen Biosystems. Altogen Biosystems shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product. This product is developed, manufactured and sold for research purposes only. Reagent is not suitable for administration to humans.