



## [Altogen Biosystems](#)

848 Rainbow Blvd Suite 823

Las Vegas, NV 89107 USA

Tel: (702) 349-6103

Fax: (702) 989-0841

Web: [www.altogen.com](http://www.altogen.com)

E-mail: [orders@altogen.com](mailto:orders@altogen.com)

### **Liver *In Vivo* Transfection Reagent (cat#5060, 5061, 5062) *In Vivo* Transfection Protocol**

Cat. No. 5060	Size: 0.5 ml
Cat. No. 5061	Size: 1.5 ml
Cat. No. 5062	Size: 8.0 ml

#### **Contents and Shipping:**

Liver *in vivo* transfection reagent is supplied in liquid form at a concentration of 1.5 mg/ml. Shipped at ambient temperature. Transfection Enhancer reagent is supplied in liquid form at a concentration of 1 mg/ml.

#### **Description:**

Liver transfection reagent is a liposome lipid-based formulation optimized for *in vivo* administration with targeted liver delivery. Liver *in vivo* transfection reagent is a proprietary, animal-origin-free formulation with high transfection efficiency of small RNA (siRNA, shRNA, microRNA) and plasmid DNA.

#### **Storage:**

Store reagent at 4°C upon receipt. If stored properly, reagent is stable for 6 months.

#### **Product Qualification:**

Reagents are tested for functional activity, absence of nuclease contamination and microbial contamination.

#### **Reagent Properties:**

Lipid conjugated complexes are stable in serum for at least 16 hours.  
Efficient delivery to the liver tissue and liver tumors via systemic administration.  
Efficient siRNA and plasmid DNA delivery via direct intratumoral injection.  
Minimal toxicity.

#### **Recommended Modes of Administration:**

Tail vein systemic intravenous (i.v.) injection.  
Direct intratumoral (i.t.) injection.  
Intraperitoneal (i.p.) injection.

#### **Intended Use:**

For research use only. Not intended for any diagnostic or therapeutic use.

#### **Safety Data Sheet (SDS):**

SDS documents are available online at <https://altogen.com>

#### **General Guidelines:**

Handle animals and conduct experiments according to national regulations and approvals by the local IACUC and ethics committee. Please note that mouse tail vein injection may require skills, experience, and associated equipment (e.g. restraining device). All persons handling animals should be properly trained.

## Guidelines for Small RNA (siRNA, shRNA, microRNA) Administration:

Prepare siRNA stock solution in DNase- and RNase-free water.

Recommended dose of siRNA (per injection):

100-120 µg siRNA with chemical modification (resistant to nuclease degradation *in vivo*)

300-500 µg standard, non-modified siRNA

## Recommended Protocol for Systemic (iv) and Intraperitoneal (ip) Administration (for 1 injection):

1. Dilute 60 µg of plasmid DNA and/or 120 µg of siRNA in 100 µl RNase-/DNase-free water. Vortex gently.
2. Add 100 µl of diluted DNA (and/or siRNA) to the sterile tube containing 50 µl Transfection Reagent.
3. Incubate for 15-20 minutes at room temperature.
4. Add 10 µl of Transfection Enhancer Reagent. Vortex gently.
5. Incubate for 5 minutes at room temperature.
6. Add required amount of sterile solution of 5% glucose (w/v):

Animal body weight (g)	Final injection volume (ml)
10-20	0.35
15-19	0.45
20-24	0.50
25-29	0.65
30-35	0.80

7. Inject animals. Please note that delivery efficiency can be significantly increased by performing secondary injection (at least 12 hours after first injection).
8. Assay for target gene expression. Maximum mRNA expression effect is typically observed 12-36 hours after injection, while maximum effect on protein level is achieved 24-48 hours post-injection.

## Recommended Protocol for Direct Intratumoral (it) Administration (for 1 injection):

1. Dilute 60 µg of plasmid DNA and/or 120 µg of siRNA in 50 µl RNase-/DNase-free water. Vortex gently.
2. Add 50 µl of diluted DNA (and/or siRNA) to the sterile tube containing 50 µl Transfection Reagent.
3. Incubate for 15-20 minutes at room temperature.
4. Add 10 µl of Transfection Enhancer Reagent. Vortex gently.
5. Incubate for 5 minutes at room temperature.
6. Inject animals.
7. Assay for target gene expression. Maximum mRNA expression effect is typically observed 12-36 hours after injection, while maximum effect on protein level is achieved 24-48 hours post-injection.

## Dosing:

Animal body weight (g)	Transfection Reagent (µl) / per Injection
10 – 35 g (mouse)	50 µl
35 – 200 g (small rodents)	100 µl
200 – 450 g (rat)	250 µl

## Limited Use Label License:

The purchase of this product conveys to the purchaser the limited right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. This product is for research purposes only and is not for use in commercial applications of any kind. For information on obtaining additional rights, please contact Altogen Biosystems at [orders@altogen.com](mailto:orders@altogen.com)

## Limited Product Warranty:

This warranty limits our liability to replacement of this product. No other warranties of any kind, express or implied, including, without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Altogen Biosystems. Altogen Biosystems shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product. This product is developed, manufactured and sold for research purposes only. Reagent is not suitable for administration to humans.